

Post-Operative Instructions
for patients of Justin T. Newman, M.D

Elbow Surgery

- Diet
 - Start with clear liquids and light foods (jello, soup, etc)
 - Progress to your normal diet as tolerated by nausea
- Wound Care
 - Keep operative dressings in place. If there is swelling of the elbow, wrist or hand – loosen the bandage about the wrist.
 - Keep your splint clean and dry.
- Medications
 - If placed, nerve blocks and pain medication injected into the wound will wear off from 8-24 hours after surgery.
 - Dr. Newman prefers that you avoid anti-inflammatory medications for 6 weeks after surgery (i.e. Advil, ibuprofen, Aleve). Occasional use is permissible, however please limit the amount taken. If they are appropriate for your surgery or condition, Dr. Newman will let you know.
 - Narcotic pain medications are often prescribed to aide in controlling your post-operative pain.
 - Length of use is generally 2-5 days, depending on level of pain
 - These medications will not alleviate all of your discomfort, but should help to manage your pain, in addition to elevation of your extremity and icing.
 - Please take medications as prescribed.
 - Do not mix with alcohol.
 - Common side effects of pain medication:
 - Nausea – take medication with food. Take anti-nausea medication if prescribed. If this persists, call the office to consider having your medication changed.
 - Drowsiness
 - Constipation – if this occurs consider taking an over-the-counter laxative such as prune juice, Senekot, Colace or Miralax
 - Do NOT drive a car or operate heaving machinery while on narcotics.
 - Pain medications will only be refilled in the post-operative period.
 - Narcotic medications cannot be called into a pharmacy. They require a written prescription from the office. Please monitor the amount of pain medications that you have in order to not run out of medications.
 - It is a strict policy that narcotics and pain medications will NOT be refilled on weekends or after hours.

- Ted Stockings
 - These compressive stockings/socks are to be worn for 2 weeks after surgery.
 - They control lower extremity swelling and decrease the chances of blood clot formation.
 - May remove for showers, but please use at all other times.
- Activity
 - OK to remove sling with minimal movement about arm for dressing, hygiene comfort
 - May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable
 - OK to use computer. Use your other hand to place your operated extremity on the desk or keyboard and allow it to fully rest there without using your own muscles to keep it in position.

EMERGENCIES

- Contact Dr. Newman or his staff if any of the following are present:
 - Painful swelling or numbness
 - An abrupt change in the nature or severity of pain
 - Fever over 101.5° (It is normal to have a low grade fever for the first day or two after surgery)
 - Redness around incisions
 - Color change in wrist, hand or lower extremity
 - Continuous drainage or bleeding from incision (routine drainage is expected)
 - Difficulty breathing
 - Excessive nausea/vomiting
 - If you have an emergency after office hours or on the weekend, call 303-344-9090 to be connected to our pager service. They will contact Dr. Newman or one of his colleagues. Do NOT call the hospital or surgery center.

**** If you have an emergency that requires immediate attention, proceed to the nearest emergency room.**

- Follow-up and questions
 - During normal business hours, Dr. Newman or his point of contact Crystal can be reached at 303-344-9090
 - If you do not already have a post-operative appointment scheduled, please call during normal business hours to schedule at 303-344-9090
 - Your first visit should be 10-14 days after surgery.
 - Plan to see Dr. Newman in clinic also at 6 weeks, 3 months and 6 months after surgery. These intervals may vary depending on your specific surgery.

Dr. Newman strives to take **exceptional** care of his patients. If there are any concerns, difficulties or problems, please let him know. He wants to work with you to achieve the best possible outcomes. This is best obtained with an open line of communication.

Wishing you a quick and successful recovery!